

66784-1

07/28/2010

1/11



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460



OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

JUL 28 2010

Subject:

EPA Registration 66784-1
Amendment Dated: June 25, 2010
EPA Received Date: June 25, 2010

Dear Ms. Olivia D. Laird

The following amendment submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA) as amended, is acceptable.

Proposed Amendment

- Revised Label per Agency letter dated April 02, 2009

General Comments

A stamped copy of the accepted label is enclosed for your records. Submit (3) copies of your final printed label before distributing or selling the product bearing the revised labeling.

Should you have any questions or comments concerning this letter, please contact Velma Noble at (703) 308-6233.


Sincerely,

A handwritten signature in black ink, appearing to read "Velma Noble".

Velma Noble
Product Manager (31)
Regulatory Management
Branch1
Antimicrobial Division (7510c)

2/11

BEFORE USING THIS PRODUCT PLEASE READ THE ENTIRE LABEL



ACTIVE INGREDIENT:	n-Alkyl (60% C ₁₄ , 30% C ₁₆ , 5% C ₁₂ , 5% C ₁₈) dimethyl benzyl ammonium chlorides	40%
INERT INGREDIENT:	Urea	60%
TOTAL		100%

This product is designed exclusively for industrial, commercial and institutional use, and must be applied by trained maintenance personnel. Label directions and precautions must be followed exactly. United Promotions Inc. will not be responsible for any injury, loss or damage if product is used in any manner not in compliance with the label directions or if the precautions are not observed.

See **DIRECTIONS FOR USE** on back of this package.

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENT
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye and skin irritation. Avoid contact with skin, eyes or clothing. Avoid inhaling or breathing dust. Wear protective eyewear, gloves, and protective clothing. Wash thoroughly with soap and water after handling and before eating, drinking or using tobacco. Remove contaminated clothing and wash before reuse.

FIRST AID

Have this product container or label with you when calling a poison control center or doctor, or going for treatment.


If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 – 20 minutes. Call a poison control center or doctor for treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15 – 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinse eye. Call a control poison center or doctor for treatment advice.

If swallowed: Call a poison center or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to unconscious person. Have person sip a glass of water if able to swallow.

If inhaled: Move person to fresh air. If person is not breathing, call 911 for an Ambulance, then give artificial respiration, preferably by mouth to mouth if possible. Call a control poison center or doctor or going for treatment.

Distributed by
United Promotions Inc.
2030 Powers Ferry Rd SE Suite 136 Atlanta, Georgia 30339 U.S.A
Telephone: 7705416055
FAX: 7705417344



Form No. 94-01

Personal Protective Equipment (PPE).

Applicators and other handlers who handle this pesticide must wear: Long sleeved shirts and long pants, socks, shoes, waterproof gloves, protective eyewear. Remove contaminated clothing and wash before reuse. Follow manufacturer's instructions for cleaning/maintaining PPE. If no instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Physical and Chemical Hazards

Do not use or store near heat or open flame. Do not mix with oxidizers, anionic soaps, and detergents.

NET WEIGHT: LBS. (KGS.)

- 2.2 lbs/1 KG
- 33 lbs/15 KG
- 300 lbs/136 KG
- 330 lbs/150 KG

EPA Reg. No. 66784-1

EPA Est. No. 66784-GA-001

STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL

ENVIRONMENTAL HAZARD

This product is toxic to fish, fresh water invertebrates, oysters, and shrimp. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of EPA.

PESTICIDE STORAGE

Store in a dry place no higher than 120°F.
Avoid creasing or impacting of sidewalls.

CONTAINER DISPOSAL

Nonrefillable container. Do not use or refill this container. Clean container promptly after emptying. Then offer for recycling or reconditioning if available, or dispose in sanitary landfill or other procedures approved by state and local authorities.

PESTICIDE DISPOSAL

Improper disposal of excess pesticides solutions or rinsate is a violation of Federal law. Waste resulting from the use of this product must be disposed of on site or at an approved waste disposal facility."

PHYSICAL AND CHEMICAL

Do not use or store near heat or open flame. Do not mix with oxidizers, anionic soaps and detergents.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its label.

DILUTION TABLE

Preparing solutions is made easy with the provided measure scoop. It holds 3.8 grams of [redacted] which is the unit or "dose" for making up solutions quickly and accurately. Add [redacted] to water. It dissolves rapidly in hot or cold water. [redacted] is stable in solution in a closed container.

YIELD/POUND OF PRONTECH™	
1600 ppm	30 gallons
800 ppm	60 gallons
400 ppm	120 gallon
200 ppm	241 gallons
100 ppm	482 gallons

ADD 1 DOSE (3.8 grams) TO:

1 qt. water for 1600 ppm
2 qts. water for 800 ppm
1 gal. water for 400 ppm
2 gals. water for 200 ppm
4 gals. water for 100 ppm

28.35 g = 1 ounce.

IN FOOD, DAIRY, MEAT, POULTRY, FISH INDUSTRIES

This product is intended for use in dairies, food processing plants, retail food markets, food service companies (cafeterias and restaurants), meat/poultry packaging and processing plants, and fish processing plants. Apply [redacted] solutions with a clean cloth, sponge, mop, or mechanical spray. [redacted] solutions are stable. Clean, unused portions may be stored in closed containers for later use. Do not contaminate feed, food, drinking water or packaging materials.

[redacted] is accepted by USDA/FSIS for use in official establishments operating under the Federal meat, poultry, shell egg grading and egg products inspecting inspection programs.

[redacted] fulfills the criteria of Appendix F, the Grade "A" Pasteurized Milk Ordinance, 1978 Recommendation of the United States Public Health Service. When tested by the method outline by Chambers, 200 ppm of the active ingredient will reduce 99.999% of the test organisms within 30 seconds, in water hardness up to 550 ppm when calculated as CaCO₃.

1. TO SANITIZE FOOD CONTACT SURFACES: Prior to sanitizing, remove gross food particles and soil by a preflush or prescrape and when necessary, a presoak. Then thoroughly wash or flush objects with a good detergent or compatible cleaner, followed by a potable water rinse.

To Sanitize Immobile Items: (tanks, chopping blocks, tables, countertops, food processing equipment): Apply a 200 ppm solution of [redacted] (3.8 grams in 2 gallons of water) to pre-cleaned, hard, nonporous immobile surfaces by thoroughly wetting surfaces with a cloth, mop, sponge, or mechanical sprayer. Surfaces must remain wet for 2 minutes. Remove excess, drain the use solution from the surface and air dry. No potable water rinse is allowed.

To Sanitize Mobile Items: (drinking glasses, dishes, eating utensils): Immerse precleaned mobile items in a 200 ppm solution of [redacted] (3.8 gms in 2 gallons of water) for 2 minutes. Remove items, drain the use solution from the surface and air dry. Prepare a fresh solution at least daily or more frequently as solution becomes dirty or soiled. No potable water rinse is allowed.

2. TO DISINFECT NON-FOOD CONTACT SURFACES Food products and packaging materials must be removed from the room or be carefully protected. Preclean heavily soiled surfaces. Apply 400 ppm solution of [redacted] to previously cleaned nonfood contact surfaces (walls, floors, counters, columns, ceilings, sinks, trash cans/barrels, light fixtures, nonfood processing equipment) with cloth, sponge, mop, or mechanical spray. Thoroughly we surfaces for 10 minutes. Allow surfaces to air dry. Prepare fresh solution daily when it becomes visibly soiled.

3. SHELL EGG SANITIZING: To sanitize clean, whole shell eggs indented for food or food product. Spray with a solution of 3.8 grams of product in 2 gallons of water (providing 200ppm active). The solution must be equal to or warmer to the eggs, but not to exceed 130°F. Wet eggs thoroughly and allow to drain. Eggs that have been sanitized with quaternary ammonium compound shall be subjected to a thorough potable water rinse only if there are to be immediately broken for use on the manufacture of egg products. Eggs must be reasonable dry before casing or breaking the solution must not be reuse for sanitizing eggs. Dirty, cracked or punctured eggs cannot be sanitized.

4. TO SANITIZE BOOTS: Shoe baths containing one inch of freshly made sanitizing solution must be placed at all entrances to buildings, production, and packaging rooms. Scrape waterproof boots. Place boots in one inch of a 200 ppm solution of [redacted] (3.8 grams in 2 gallons of water) for 2 minutes before entering building, production, and packaging room. Change sanitizing solution in bath at least daily or sooner if solution appears dirty or soiled.

5. GLOVE DIP SANITIZING: Dip prewashed (plastic, latex or other synthetic rubber) non-porous gloved hands into a suitable clean container that holds one inch of a 220 ppm solution of [redacted] (3.8 gms in 2 gallons of water) sanitizing solution for 2 minutes. Change the sanitizing solution in the bath at least daily or when solution appears dirty or soiled."

ANIMAL PREMISES

6. FARMS

Remove all animals and feed from premises, vehicles, and enclosures. Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes, and other facilities and fixtures occupied or traversed by animals. Empty all troughs, racks, and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate all surfaces with a use solution of 400 ppm of this product for a period of 10 minutes. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure. Ventilate buildings, cars, boats, and other closed spaces. Do not house livestock or employ equipment until treatment has been absorbed, set, or dried. Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains, and waters with soap or detergent, and rinse with potable water before reuse.

7. POULTRY FARMS

Use it to disinfect poultry hatcheries, setter trays, hatch baskets and machinery, all dry storage, cold rooms, docks, trucks, and transport cages.

MICROORGANISM*	TYPE OF MICROORGANISM
<i>Avian influenza H9N9</i>	Virus
<i>Avian influenza H5N1</i>	Virus
<i>Swine Flu H1N1</i>	Virus
<i>Aujesky disease virus</i>	Virus
<i>Avian metapneumovirus</i>	Virus
<i>Gumboro virus</i>	Virus

Remove all poultry and feeds from premises, trucks, coops, and crates. Remove all litter and droppings from floors, walls and surfaces of facilities occupied or traversed by poultry. Empty all troughs, racks, and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate surfaces with a use solution of 400 ppm of this product for a period of 10 minutes. Ventilate buildings, coops, and other closed spaces. Do not house poultry or employ equipment until treatment has been absorbed, set, or dried. Thoroughly scrub treated feed racks, troughs, automatic feeders, fountains, with soap or detergent, and rinse with potable water before reuse.

8. VETERINARY FACILITIES: For cleaning and disinfecting the following hard non-porous surfaces: instruments, cages, kennels, stables, catteries, etc. Remove all animals and feed from premise, animal transportation vehicles, crates, etc. Remove all litter, droppings and manure from floors, walls and surfaces occupied or transverse by animals. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate surfaces with a use solution of 400 ppm of this product for a period of 10 minutes. Ventilate building and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried. Thoroughly scrub all treated feed racks, automatic feeders, waterers, and other equipment which dispenses food or water with soap or detergent, and rinse with potable water before reuse.

PETS

1. DIRECTIONS FOR DISINFECTION AND ODOR CONTROL IN HOUSEHOLDS: For cleaning/disinfecting the following hard non-porous surfaces: carpets, floors, furniture, cages, litter boxes, bird cages and pet living and sleeping areas. Remove all litter, droppings and manure from floors, walls and surfaces occupied or transverse by animals. Thoroughly clean all surfaces with soap or detergent and rinse with

water. Apply a fine spray mist of 800 ppm of this product throughout the area, and it will effectively destroy/eliminate offensive odors, caused by bacteria, on contact to break the re-soiling cycle.

Bacteria	Viruses	Fungi
<i>Staphylococcus aureus</i> (Staph)	<i>Rotavirus</i> (the leading cause of infectious diarrhea in children)	<i>Trichophyton mentagrophytes</i> (the fungus that causes athlete's foot)
<i>Salmonella enterica</i>	<i>Influenza A</i> (flu)	<i>Aspergillus</i>
<i>Salmonella thyphimurium</i>	<i>Avian Influenza A (H5N1, H9N9)</i>	
<i>Listeria monocytogenes</i> (Listeria)	<i>HIV-1</i>	
<i>Enterococcus faecalis</i>		
<i>Escherichia coli</i> (E.coli)		
<i>Campylobacter jejuni</i>		
<i>Pseudomonas aeruginosa</i>		
<i>Bacillus cereus</i>		
<i>Klebsiella pneumoniae</i>		
<i>Streptococcus agalactiae</i>		
<i>Proteus vulgaris</i>		
<i>Haemophilus influenza</i>		
<i>Pasteurella multocida</i>		

Bacteria	Viruses	Fungi
<i>Staphylococcus aureus</i> (Staph)	<i>Rotavirus</i> (the leading cause of infectious diarrhea in children)	<i>Trichophyton mentagrophytes</i> (the fungus that causes athlete's foot)
<i>Salmonella enterica</i>	<i>Influenza A</i> (flu)	<i>Aspergillus</i>
<i>Salmonella thyphimurium</i>	<i>Avian Influenza A (H5N1, H9N9)</i>	
<i>Listeria monocytogenes</i> (Listeria)	<i>HIV-1</i>	
<i>Enterococcus faecalis</i>		
<i>Escherichia coli</i> (E.coli)		
<i>Campylobacter jejuni</i>		
<i>Pseudomonas aeruginosa</i>		
<i>Bacillus cereus</i>		
<i>Klebsiella pneumoniae</i>		
<i>Streptococcus agalactiae</i>		
<i>Proteus vulgaris</i>		
<i>Haemophilus influenza</i>		
<i>Pasteurella multocida</i>		

GENERAL USES

1. **TO CONTROL ODORS** in dry storage areas, cold rooms, refrigerated showcases, kitchens, docks, dumpsters, and in waiting rooms, toilet exteriors, showers, etc., apply a fine spray mist of 800 ppm solution throughout the area or on the equipment. To control more intense odors, use a 1600 ppm solution. **Rinse food contact surfaces with potable water before use.** Do not spray on food or packaging materials.

2. **GLOVE DIP SANITIZING:** Dip prewashed (plastic, latex or other synthetic rubber) non-porous gloved hands into a suitable clean container that holds one inch of a 220 ppm solution of (3.8 gms in 2 gallons of water) sanitizing solution for 2 minutes. Change the sanitizing solution in the bath at least daily or when solution appears dirty or soiled.

PULP & PAPER

is included in the list of substances acceptable as slimeicides in 21 CFR 176.300 (c) without limitations and (d). may be used as an antimicrobial agent to control slime in the manufacturing of paper and paper board, including those paper products that contact food.

7/11

AS A SLIMICIDE IN PULP AND PAPER MILLS: To control bacteria and fungi in the process water, add 340 grams of [redacted] to each 1,000 gallons of water. Dilution will be 36 ppm. Determine results and adjust the amount of [redacted] to the minimum necessary for control.

Add [redacted] once a day to any or several of the following areas as needed: pulper, machine chest, fan pump suction, mix chest, broke chest.

AS AN ALGICIDE IN PULP AND PAPER MILLS: To control algae in process water, add 57 to 113 grams of [redacted] to each 1,000 gallons of water, depending on the kind and amount of algae present. Dilutions will be 6 ppm to 12 ppm.

Add [redacted] once a day to any or several of the following areas as needed: pulper, machine chest, fan pump suction, mix chest, broke chest.

RECIRCULATING COOLING TOWERS SYSTEMS

FOR ALGAE CONTROL IN INDUSTRIAL AND COMMERCIAL RECIRCULATING COOLING TOWERS SYSTEMS:

To control algae in metal or wooden cooling towers, apply the product to the sources of water such as the boiler and cooling tower. Apply after cleaning the system. Apply directly to the water of the boiler and cooling tower, following modified intermittent parent method. Initial dose:

3.8 grams of [redacted] in 4 gallons water = 100 ppm
1 ounce of [redacted] in 30 gallons water = 100 ppm

When control is achieved maintain with 20 ppm applications weekly.

FOR ALGAE CONTROL IN INDUSTRIES AND COMMERCIAL WATER COOLING ONCE THROUGH SYSTEMS:

To control algae in metal or wooden recirculating water cooling systems, apply the product to the sources of water such as the reservoir. Apply after cleaning the system. Apply directly to the water of the reservoir, following modified intermittent parent method. Initial dose:

3.8 grams of [redacted] in 4 gallons water = 100 ppm
1 ounce of [redacted] in 30 gallons water = 100 ppm

When control is achieved maintain with 20 ppm applications weekly.

DO NOT APPLY THIS PRODUCT MORE THAN 4 TIMES PER YEAR.

Deactivation: This product must be deactivated prior to discharge of the NPDES outfall.

To Deactivate: Use Bentonite Clay at a minimum ratio 5 ppm clay to 1 ppm product. Deactivation must occur prior to discharge of the NPDES outfall.

SWIMMING POOLS

AS A SWIMMING POOL ALGICIDE:

Keeps pool water clear and free of visible algae.

*Initial Dose Ratio: 907 grams in 50,000 gallons of water
*Maintenance Dose Ratio: 227 grams in 50,000 gallons of water

When used as directed, this product will kill and prevent development of *Chlorella pyrenoidosa* (green algae), *Peridinium inondetum* (black algae), *Phormidium retzii* (Square D Algae), *Lyngbya versicolor* (blue-green algae) and other swimming pool algae. It is compatible with most commonly used swimming pool chemicals, but must not be mixed with soap or other water treating chemicals. It works best when added by itself. Add [redacted] directly into the skimmer or vacuum line to the filter, recirculating the filter and returning the compound in the filter to the pool. Keep the filter running for 12 hours. Backwash the filter if it becomes clogged. Do not apply when swimmers are in the pool. 15 minute Re-entry interval for swimmers.

INITIAL DOSE is used when filling the pool or at the start of the swimming season if there is visible algae growth. Add [redacted] when the filter and pump are in operation for better mixing and distribution. Vacuum the pool after 24 hours to remove any algae debris. If visible algae are still present or reappear, repeat treatment procedures as necessary. When pool is free of visible algae, use maintenance dose.

MAINTENANCE DOSE is used at 3 - 5 day intervals. Add [redacted] while water is being circulated. Use at 3-day intervals when temperatures are high, when the bathing load is heavy and always after a rain. Also add a maintenance dose for each 10,000 gallons of makeup water or in that ratio. Add 45 grams for 10,000 gallons, 113 grams for 25,000 gallons, etc. After adding [redacted], wait 15 minutes before entering pool.

HOSPITAL USE DIRECTIONS

For cleaning and disinfecting the following hard non-porous surfaces: Preclean heavily soiled surfaces. Apply 800 ppm solution of [redacted] to previously cleaned nonfood contact surfaces (walls, floors, bedframes, tables, counters, columns, ceilings, sinks,, trash cans/barrels, light fixtures) with cloth, sponge, mop, or mechanical spray. Thoroughly wet surfaces for 10 minutes. Allow surfaces to air dry. Prepare fresh solution daily or when it becomes visibly soiled.

*HIV-1

KILLS HIV ON PRECLEANED ENVIRONMENTAL SURFACES/OBJECT PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood with body fluids and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission human immunodeficiency virus Type 1 (HIV-1) (associated with AIDS).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 ON SURFACES/OBJECTS SOILED WITH BLOOD OR BODY FLUIDS.

At 3.8 grams per 2 gallons of water for a 4-minute contact time this product was found to be effective against HIV-1 (AIDS Virus).

PERSONAL [redacted]: Specific barrier protection items to be used when handling items soiled with blood and body fluid are disposable latex glove, gowns, masks, and eye coverings.


CLEANING PROCEDURE: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product.

DISPOSAL OF INFECTIOUS MATERIAL: Blood and other body fluids must be autoclaved and disposed of according to federal, state and local regulations for infectious waste disposal.

CONTACT TIME: Leave surface wet for 4 minutes for HIV-1. This 4 minute contact time will not control other organisms listed on this label

MOLD AND MILDEW

Add 3.8 grams per 2 gallons of water to control the growth of mold and mildew and their odors on hard, non-porous surfaces. Thoroughly wet all treated surfaces completely. Let air dry. Repeat application weekly or when growth or odor reappears.



9/11

FLORICULTURE

See Statement of Practical Treatment on Back Panel

Net weight - 2.2 lbs/1 KG
 - 33 lbs/15 KG
 - 300 lbs/136 KG
 - 330 lbs/150 KG

User Safety Requirements

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove PPE immediately after handling this product. Wash outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing

Fungicide - Bactericide - Algacide

For control of foliar and root diseases in Floriculture crops in greenhouses, shadehouses, Garden Centers, Lanscapes, Nurseries, Golf Courses.

Do not apply this product in a way that will contact workers or other persons either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation

Personal Protection Equipment (PPE) Applicators and other handlers who handle this pesticide must wear: Coveralls or long sleeved shirt, long pants, socks, chemical resistant footwear and gloves. Also wear protective eyewear such as goggles or face shield, dust mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C) or respirator with an organic vapor removing cartridge with a prefilter approved for pesticides (MSH/NIOSH approval number TC23-C).

Discard clothing and other absorbant materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no instructions or washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Agricultural Use Requirements: Use this product only in accordance with it's labeling and with the Worker Protection Standard, **40 CFR part 170**. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into the treated areas during the restricted entry interval (REI) of 48 hours.

Application is limited to spot treatment of diseased ornamental plants and flowers at a maximum of 800ppm and total use of 5lbs/ A.I acre

ORNAMENTAL CROPS	DISEASE	INTERVAL OF APPLICATION	TYPE OF APPLICATION	RATE (PPM*)
Ornamental Crops	Disease	Interval of Application	Type of Application	Rate (PPM)
Such as				*parts per million
Carnation	Fusarium oxysporum	15 days	Drench	1600
(Dianthus caryophyllus)	Fusarium roseum	5 days	Spray	800
	Botrytis cinerea	8 days	Spray	200-400
	Cladosporium echinulatum	8 days	Spray	400
	Alternaria sp.	8 days	Spray	800
	Pythium sp.	15 days	Drench	800
	Rhizoctonia sp.	15 days	Drench	800
Rose	Botrytis cinerea	8 days	Spray	200-400
	Rhizoctonia sp.	15 days	Drench	800
	Oidium sp. (Powdery Mildew)	4-8 days	Spray	400
	Agrobacterium tumefaciens	1 time	Drench	800

	Pythium	15 days	Drench	800
Chrysanthemum (Chrysanthemum morifolium)	Botrytis cinerea	8 days	Spray	200-400
	Alternaria sp.	8 days	Spray	600-800
	Pythium sp.	15 days	Drench	800
	Pseudomonas cichorii	15 days	Spray	400-600
Baby's Breath (Gypsophila paniculata)	Botrytis cinerea	8 days	Spray	200-400
	Agrobacterium tumefaciens	1 time	Drench	800
	Erwinia carotovora	15 days	Drench	800
Statice (Limonium sp.)	Botrytis cinerea	8 days	Spray	200-400
	Pythium sp.	15 days	Drench	800
	Rhizoctonia sp.	15 days	Drench	800
Aster (Callistephus chinensis)	Fusarium roseum	8 days	Spray	800
	Botrytis cinerea	8 days	Spray	200-400
	Oidium sp. (Powdery Mildew)	4-8 days	Spray	400
	Agrobacterium tumefaciens	1 time	Drench	800
Gerbera (Gerbera jamesonii)	Fusarium roseum	8 days	Spray	800
	Botrytis cinerea	8 days	Spray	200-400
	Pythium sp.	15 days	Drench	800
	Rhizoctonia	15 days	Drench	800
	Pseudomonas cichorii	15 days	Spray	400-600
	Xanthomonas sp.	15 days	Spray	400-600
Alstroemeria (Alstroemeria sp.)	Botrytis cinerea	8 days	Spray	200-400
	Pythium sp.	15 days	Drench	800
	Rhizoctonia sp.	15 days	Drench	800

ORNAMENTAL CROPS Such As	DISEASE	INTERVAL OF APPLICATION	TYPE OF APPLICATION	RATE (PPM)
Snapdragon (Antirrhinum majus)	Botrytis cinerea	8 days	Spray	200-400
	Alternaria sp.	8 days	Spray	800
	Pythium sp.	15 days	Drench	800
	Rhizoctonia sp.	15 days	Drench	800
Delphinium sp. (Delphinium sp.)	Botrytis cinerea	8 days	Spray	200-400
	Alternaria sp.	8 days	Spray	800
	Pythium sp.	15 days	Drench	800
	Rhizoctonia sp.	15 days	Drench	800
Trachelium (Trachelium caeruleum)	Botrytis cinerea	8 days	Spray	200-400
	Alternaria sp.	8 days	Spray	800
	Pythium sp.	15 days	Drench	800
	Rhizoctonia sp.	15 days	Drench	800
Stocks (Matthiola sp.)	Botrytis cinerea	8 days	Spray	200-400
	Alternaria sp.	8 days	Spray	800
	Pythium sp.	15 days	Drench	800
	Rhizoctonia	15 days	Drench	800
Anthurium (Anthurium sp.)	Xanthomonas campestris	15 days	Spray	400-600
	Xanthomonas anthuriae	7 days	Spray	400-600
Post-Harvest Cut Flowers	Bacteria, Fungi	1 time	Immerse stems 1-2 hours	200
Turfgrass Grass or Dichondra.	Rhizoctonia sp.	10-14 days	Spray	800-1600
	Phyium sp.	10-14 days	Spray	800-1600
	Xanthomonas sp.	10-14 days	Spray	400-600
	Pseudomonas sp.	10-14 days	Spray	400-600
	Pyricularia oryzae	10-14 days	Spray	800
	Sclerotinia homeocarpa	10-14 days	Spray	800-1600
	Helminthosporium	10-14 days	Spray	400-600
	Marasmius spp	3 days	Spray	1600
	Lycoperdon spp	3 days	Spray	1600
	Agrocybe pedates	3 days	Spray	1600
	Bovistia plumbea	3 days	Spray	1600
	Fusarium blight	3 days	Spray	1600

Preventive Rate: 1/2 oz per 1000 sq. ft. in 2 gallons of water.
Curative Rate: 1 oz per 1000 sq. ft. in 2 gallons of water.

NOTE:

1. Using less carrier may not produce the desired results
2. When using a defoamer, use the lowest labeled rate.

Disinfestation (Floriculture species including those listed above)	Bacteria, Fungi	1 time	Immerse in solution 30 seconds	400
--	-----------------	--------	-----------------------------------	-----

TO DISINFECT HARD NON – POROUS SURFACES

Remove gross filth and heavy soil with a preliminary cleaning step by flushing with water, vacuuming or sweeping. Use [REDACTED] to disinfect any previously cleaned and rinsed surface (e.g., floors, walls, benches, columns, ceilings, light fixtures, sinks, trash barrels, inedible drums, evaporative coolers, equipment) apply a 400 ppm solution as needed, of [REDACTED] (3.8 grams in one gallon of water equals 400 ppm) using a brush or mechanical spray device. Wet surfaces thoroughly and allow 10 minutes contact time. Do not rinse unless required by local authorities. This application is also suitable for toilets exteriors, showers, locker rooms, break rooms and similar public areas. To disinfect equipment and tools immerse them daily in a 400 ppm solution for 10 minutes, of [REDACTED] (3.8 grams in one gallon of water equals 400 ppm). Prepare fresh use solution daily. Do not reuse solution. Discard solutions when they become soiled.

MIXING AND APPLICATION INSTRUCTIONS

Mix [REDACTED] with water according to the rates listed in Directions For Use and the [REDACTED] Dilution Table. Spray Floriculture crops to run off on upper and lower leaf surfaces, using 50 to 100 gallons of solution per acre. When applying [REDACTED] as a drench, mix [REDACTED] with water at the correct rate listed under Directions For Use, and drench at the rate of 100 gallons per 400 square feet, or apply in sufficient amount to saturate the soil mixture.

When spraying [REDACTED] in greenhouses or shadehouses, make sure vents or sidewalls are open during and immediately after application to supply adequate ventilation.

See Front Panel for
Precautionary Statement

Refer to DILUTION TABLE
on side panel.

Do not mix with soap anionic detergents, which would reduce the effectiveness of [REDACTED]

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants (a) that this product conforms to the chemical description on the label; (b) that this product is reasonable fit for the purposes set forth in the directions for use when it is used in accordance with such directions; and (c) that the directions, warnings and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants. Tests have not been made on all varieties or in all states or under all conditions.

THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESSED OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS.

EPA Reg. No. 66784-1
Form No. 94-03P